


# Variation of ice crystal shape, size & asymmetry parameter in tropical deep convection retrieved from combined MODIS and POLDER measurements

Bastiaan van Diedenhoven

(Columbia University, NASA GISS)

Brian Cairns, Ann Fridlind and Andrew Ackerman


(NASA GISS)



National Aeronautics and Space Administration  
**Goddard Institute for Space Studies**  
New York, N.Y. 10025


# Current knowledge on global ice cloud properties

e) Cloud Optical Thickness Ice



Maddux et al.,  
JAOT 2010

c) Cloud Effective Radius Ice ( $\mu\text{m}$ )



## Ice cloud asymmetry parameter:

Models: 0.6-0.95

In situ:  $\sim 0.75$


?

?

Required accuracy: 0.01-0.04 (Vogelmann & Ackerman, 1995)

# Ice crystal asymmetry parameter

- Ice crystal asymmetry parameter mainly depends on
  - Shape (or shape of components)
  - Distortion/Microscopical Roughness/impurity
- Asymmetry parameter of aggregates similar to components



# Information from Polarization

Polarization contains info about


- Aspect ratio
- Roughness

Multi-directional polarized  
reflectance measurements

conserve

Single scattering features

Retrieve aspect ratio and  
roughness from polarization to  
estimate asymmetry parameter



# Retrieval procedure

**LUT of polarized reflectances for columns/plates**


51 aspect ratios 0.02 – 50

15 roughness values ( $\delta = 0 - 0.7$ )

**Find best fit to measured multi-directional polarized reflectances**

At least scattering angles  $120^\circ - 150^\circ$

**Asymmetry parameter of best-fit hexagon is retrieved value**



**Use retrieved asymmetry parameter for cloud optical thickness retrieval (later also effective radius)**


# Simulated data test

## Simulated data:

- Complex ice habits (Yang et al.)
- 3 roughness degrees
- 20 different size distributions


## Retrieved asymmetry parameter

- Within 5% (0.04)
- Mean bias: 0.004
- Standard deviation: 0.02





Van Diedenhoven et al.,  
Atmos. Meas. Tech., 5, 2361–2374, 2012

# Application to POLDER/PARASOL and MODIS/Aqua




# TWP-ICE campaign

- ARM/NASA campaign in early 2006
- Sampling Asian monsoon in Northern Australia
- Here MODIS/POLDER data collected for extended period: 16 Jan.-20 Feb.




# POLDER retrieval statistics COT>5

- ONLY retrievals for COT>5
- Asymmetry parameter
  - $\sim 0.75-0.78$
- Aspect ratio
  - $\sim 0.4-0.7$
  - Mostly plate-like
- Distortion
  - $\sim 0.4-0.7$  (rough-severely rough)




# A more detailed look at statistics


2006/01/16 – 2006/01/23




2006/01/25 – 2006/02/13



2006/02/14 – 2006/02/20






Rainfall




# Vertical profiles

- Median values and IQR
- With increasing height
  - decreasing  $g$
  - increasing aspect ratio
  - increasing distortion
- Profiles depend on atmospheric condition




# Vertical profiles $g$ and $R_{\text{eff}}$

- With increasing height
  - decreasing  $g$
  - decreasing  $R_{\text{eff}}$
- Profiles depend on atmospheric condition



- C5  $R_{\text{eff}}$  biased by 1-5 micron due to high  $g$  assumed!
- Part of  $R_{\text{eff}}$  profile could explained by  $g$  profile






# TWP-ICE, TC4 & CRYSTAL-FACE

- Deep convection ( $\text{COT} > 5$ )
  - $g \sim 0.75-0.79$
  - Aspect ratios  $\sim 0.2-1.2$
  - Distortion  $\sim 0.4-0.7$

From 4 case studies using airborne RSP!

Van Diedenhoven et al. ACP, 2013







# Future work

- POLDER retrievals for unsaturated polarized reflection (COT<5)
  - Proper surface
  - Rayleigh
- Effective radius retrievals based on retrieved asymmetry parameters
- Speed up algorithm
- Global MODIS/POLDER retrievals
- Validation studies using new processing techniques for in situ CPI images (Greg McFarquhar)

# Caution! Thick cloud bias?

- Previous global POLDER analysis
  - show narrow range of polarized reflectance
  - assume saturation of polarized reflectance
- But  $R_p$  only saturates at  $COT \sim 5$
- Where are all the cloud  $COT < 5$  ??

Polarized reflectance at scattering angle  $120^\circ$  for specified COT vs saturated value



**DRYDEN**  
FLIGHT RESEARCH CENTER




L. PU  
Z. PU

# Backup slides

# Shortwave flux


- Cloud optical thickness =4
- Cloud between 10-11 km
- SZA=60 degrees



From “An accurate and flexible parameterization for shortwave optical properties of ice crystals”, van Diedenhoven et al., in preparation


# POLDER vs MODIS C5 optical thickness

- MODIS C5 biased high by  $\sim 0.83$ 
  - Makes sense:  
 $(1-0.77)/ (1-0.82) = 0.78$
- Thick clouds have stronger bias!
- Similar to results of Zhang et al. 2009




# POLDER vs MODIS optical thickness


- Zhang et al.  
2009 compares  
POLDER  
level-2 product  
to MODIS C5
- Level-2 uses  
 $g=0.74$ 
  - $(1-0.74)/$   
 $(1-0.82)=$   
0.69




# Retrievals mean




# Retrievals standard deviation







# POLDER Liquid index

- Fit straight line through  $120^\circ$ - $160^\circ$  measurements
- Ice index =  
 $\text{Mean}(|\text{fit-measurement}|)$
- Straight-forward to simulate from model




# Physical interpretation of liquid index

- Liquid index
  - Indicates to what degree liquid is *obscured* by ice above
  - $\sim 2\text{-}3$  for pure water clouds
  - $\sim 0$  for pure ice clouds or ice *topped* clouds



# Determining Aspect Ratios from CPI (McFarquhar et al.)



# Determining Aspect Ratios from CPI (McFarquhar et al.)

